


# PRV

PATENT- OCH REGISTRERINGSVERKET  
Patentavdelningen

SE 99 / 01075

PCT/ SE 99 / 01075  
9 / 555575



## Intyg Certificate

5



Härmed intygas att bifogade kopior överensstämmer med de handlingar som ursprungligen ingivits till Patent- och registreringsverket i nedannämnda ansökan.

Ansökan ingavs ursprungligen på engelska.

This is to certify that the annexed is a true copy of the documents as originally filed with the Patent- and Registration Office in connection with the following patent application.

The application was originally filed in English.


(71) Sökande Astra Pharma Inc, Mississauga CA  
Applicant (s)

(21) Patentansökningsnummer 9802209-8  
Patent application number

(86) Ingivningsdatum 1998-06-22  
Date of filing

Stockholm, 1999-07-22

För Patent- och registreringsverket  
For the Patent- and Registration Office

  
Evy Morin

Avgift  
Fee

**PRIORITY DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH  
RULE 17.1(a) OR (b)

PATENT- OCH  
REGISTRERINGSVERKET  
SWEDEN

Postadress/Adress  
Box 5055  
S-102 42 STOCKHOLM

Telefon/Phone  
+46 8 782 25 00  
Vx 08-782 25 00

Telex  
17978  
PATOREG S

Telefax  
+46 8 666 02 86  
08-666 02 86

## NOVEL COMPOUNDS

Field of the invention

5 The present invention is related to novel compounds, to a process for their preparation, their use and pharmaceutical compositions comprising the novel compounds. The novel compounds are useful in therapy, and in particular for the treatment of pain.

Background and prior art

10

The  $\delta$  receptor has been identified as having a role in many bodily functions such as circulatory and pain systems. Ligands for the  $\delta$  receptor may therefore find potential use as analgesics, and/or as antihypertensive agents. Ligands for the  $\delta$  receptor have also been shown to possess immunomodulatory activities.

15

The identification of at least three different populations of opioid receptors ( $\mu$ ,  $\delta$  and  $\kappa$ ) is now well established and all three are apparent in both central and peripheral nervous systems of many species including man. Analgesia has been observed in various animal models when one or more of these receptors has been activated.

20

With few exceptions, currently available selective opioid  $\delta$  ligands are peptidic in nature and are unsuitable for administration by systemic routes. Some non-peptidic  $\delta$  antagonists have been available for some time (see Takemori and Portoghese, 1992, Ann. Rev. Pharmacol. Tox., 32: 239-269. for review). These compounds, e.g. naltrindole, suffer from  
25 rather poor (i.e., < 10-fold) selectivity for the  $\delta$  receptor vs.  $\mu$  receptor binding and exhibit no analgesic activity, a fact which underscores the need for the development of highly selective non-peptidic  $\delta$  ligands.


Thus, the problem underlying the present invention was to find new analgesics having improved analgesic effects, but also with an improved side-effect profile over current  $\mu$  agonists and potential oral efficacy.

5 Analgesics that have been identified and are existing in the prior art have many disadvantages in that they suffer from poor pharmacokinetics and are not analgesic when administered by systemic routes. Also, it has been documented that preferred compounds, described within the prior art, show significant convulsive effects when administered systemically.

10

The problem mentioned above has now been solved by developing novel 1,4-substituted phenyl compounds, as will be described below.

15 The novel compounds according to the present invention are defined by the general formula I



20 wherein

**m and n** is each and independently an integer of from 0-3, and one or more of the hydrogens in such an alkylene-chain may optionally be substituted by anyone of C<sub>1</sub> - C<sub>6</sub> alkyl, C<sub>1</sub> - C<sub>6</sub> alkoxy, or hydroxy; *or*

25 one or more of the methylene groups may optionally be substituted by a heteroatom such as O, N or S;

$R^1$  is selected from hydrogen, a branched or straight  $C_1$ - $C_6$  alkyl,  $C_1$ - $C_6$  alkenyl,  $C_3$ - $C_8$  cycloalkyl,  $C_4$ - $C_8$ (alkyl-cycloalkyl) wherein alkyl is  $C_1$ - $C_2$  alkyl and cycloalkyl is  $C_3$ - $C_6$  cycloalkyl;

$R^2$  is selected from any of

(i) hydrogen;

(ii) a straight or branched  $C_1$ - $C_6$  alkyl, said alkyl optionally comprising one or more saturated bonds (double or triple bonds) within the alkyl chain;

(iii)  $-(CH_2)_q$ - aryl];

(iv)  $-(CH_2)_r$ - heteroaryl] where the heteroaryl has from 5 to 10 atoms and the heteroatom being selected from any of S, N and O;

and wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents Y where each Y is as defined below; and wherein q and r is each and independently an integer of from 0 to 3;

(v)  $C_3$ - $C_{10}$  cycloalkyl, optionally comprising one or more unsaturations and optionally substituted by one or more heteroaryl(s) where the heteroaryl has from 5 to 10 atoms and the heteroatom being selected from any of S, N and O;

and wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents Y where each Y is as defined below;

(vi)  $C_6$ - $C_{10}$  aryl, optionally and independently substituted by one or more heteroaryl(s) having from 5 to 10 atoms and the heteroatom(s) being selected from any of S, N and O

and wherein the heteroaryl may optionally and independently be substituted by 1 or 2 substituents **Y** wherein each **Y** is as defined below;

- 5 (vii) heteroaryl having from 5 to 10 atoms and the heteroatom being selected from any of **S, N and O**; wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents **Y** wherein each **Y** is as defined below;

*or*

10

**R<sup>1</sup>** and **R<sup>2</sup>** may optionally form a heterocyclic ring;

**R<sup>3</sup>** is selected from anyone of

15

(i) hydrogen;

(ii) a straight or branched **C<sub>1</sub>-C<sub>6</sub>** alkyl, said alkyl optionally comprising one or more saturated bonds (double or triple bonds) within the alkyl chain;

20

(iii) **-(CH<sub>2</sub>)<sub>q</sub>-aryl** wherein **q** is an integer of from 0 to 3, and wherein the aryl may optionally be substituted by one or more heteroaryl(s) having from 5 to 10 atoms and the heteroatom being selected from any of **S, N and O**; and wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents **Y** wherein each **Y** is as defined below;

25

(iv) **heteoaryl-(C<sub>5</sub> - C<sub>10</sub> alkyl)**, where the heteoaryl has from 5 to 10 atoms and the heteroatom being selected from any of S, N and O, and wherein the aryl and heteoaryl may optionally and independently be substituted by 1 or 2 substituents Y where each Y is as defined below;

5

(v) **C<sub>3</sub>-C<sub>10</sub> cycloalkyl**, optionally comprising one or more unsaturations and optionally substituted by one or more heteoaryl(s) having from 5 to 10 atoms and the heteroatom being selected from any of S, N and O, and wherein the aryl and heteoaryl may optionally and independently be substituted by 1 or 2 substituents Y where each Y is as defined below;

10

(vi) **-[(C<sub>3</sub>-C<sub>6</sub> cycloalkyl)-(CH<sub>2</sub>)<sub>q</sub>]** wherein q is an integer of from 1 to 3;

**R<sup>4</sup>** is selected from

15

(i) **hydrogen**;

(ii) **a straight or branched C<sub>1</sub>-C<sub>6</sub> alkyl**, said alkyl optionally comprising one or more saturated bonds (double or triple bonds) within the alkyl chain;

20

(iii) **-[(CH<sub>2</sub>)<sub>q</sub>-aryl]** wherein q is an integer of from 0 to 3, and wherein the aryl may optionally be substituted by one or more heteoaryl(s) having from 5 to 10 atoms and the heteroatom being selected from any of S, N and O; and wherein the aryl and heteoaryl may optionally and independently be substituted by 1 or 2 substituents Y wherein each Y is as defined below;

25

(iv) **heteoaryl-(C<sub>5</sub> - C<sub>10</sub> alkyl)**, where the heteoaryl has from 5 to 10 atoms and the heteroatom being selected from any of S, N and O, and wherein the aryl and heteoaryl may

optionally and independently be substituted by 1 or 2 substituents **Y** where each **Y** is as defined below;

5 (v) **C<sub>3</sub>-C<sub>10</sub> cycloalkyl**, optionally comprising one or more unsaturations and optionally substituted by one or more heteroaryl(s) where the heteroaryl has from 5 to 10 atoms and the heteroatom being selected from any of S, N and O;

and wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents **Y** where each **Y** is as defined below;

10 (vi) **C<sub>6</sub>-C<sub>10</sub> aryl**, optionally and independently substituted by one or more heteroaryl(s) having from 5 to 10 atoms and the heteroatom(s) being selected from any of S, N and O and wherein the heteroaryl may optionally and independently be substituted by 1 or 2 substituents **Y** wherein each **Y** is as defined below;

15

(vii) **heteroaryl having from 5 to 10 atoms and the heteroatom being selected from any of S, N and O**; wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents **Y** wherein each **Y** is as defined below;

20 **R<sup>5</sup>** is selected from anyone of

(i) **hydrogen**;

25 (ii) **a straight or branched C<sub>1</sub>-C<sub>6</sub> alkyl**, said alkyl optionally comprising one or more saturated bonds (double or triple bonds) within the alkyl chain;


(iii) **-[(CH<sub>2</sub>)<sub>q</sub>-aryl]** wherein **q** is an integer of from 0 to 3, and wherein the aryl may optionally be substituted by one or more heteroaryl(s) having from 5 to 10 atoms and the heteroatom being selected from any of S, N and O; and wherein the aryl and heteroaryl

may optionally and independently be substituted by 1 or 2 substituents Y wherein each Y is as defined below;

(iv) **heteoaryl-(C<sub>5</sub> - C<sub>10</sub> alkyl)**, where the heteoaryl has from 5 to 10 atoms and the heteroatom being selected from any of S, N and O, and wherein the aryl and heteoaryl may optionally and independently be substituted by 1 or 2 substituents Y where each Y is as defined below;

(v) **C<sub>3</sub>-C<sub>10</sub> cycloalkyl**, optionally comprising one or more unsaturations and optionally substituted by one or more heteoaryl(s) having from 5 to 10 atoms and the heteroatom being selected from any of S, N and O, and wherein the aryl and heteoaryl may optionally and independently be substituted by 1 or 2 substituents Y where each Y is as defined below;

15 (vi)



wherein

20 **R<sup>7</sup>, R<sup>8</sup>, R<sup>9</sup>, R<sup>10</sup> and R<sup>11</sup>** is each and independently selected from

(a) **hydrogen;**

(b) **a straight or branched C<sub>1</sub>-C<sub>6</sub> alkyl**, said alkyl optionally comprising one or more saturated bonds (double or triple bonds) within the alkyl chain;

25



(c)  $-(\text{CH}_2)_q\text{-aryl}$ ] wherein  $q$  is an integer of from 0 to 3, and wherein the aryl may optionally be substituted by one or more heteroaryl(s) having from 5 to 10 atoms and the heteroatom being selected from any of S, N and O; and wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents **Y** wherein each **Y** is as defined below;

(d) **heteoaryl-(C<sub>5</sub> - C<sub>10</sub> alkyl)**, where the heteroaryl has from 5 to 10 atoms and the heteroatom being selected from any of S, N and O, and wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents **Y** where each **Y** is as defined below;

(e) **C<sub>3</sub>-C<sub>10</sub> cycloalkyl**, optionally comprising one or more unsaturations and optionally substituted by one or more heteroaryl(s) where the heteroaryl has from 5 to 10 atoms and the heteroatom being selected from any of S, N and O; and wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents **Y** where each **Y** is as defined below;

(f) **C<sub>6</sub>-C<sub>10</sub> aryl**, optionally and independently substituted by one or more heteroaryl(s) having from 5 to 10 atoms and the heteroatom(s) being selected from any of S, N and O and wherein the heteroaryl may optionally and independently be substituted by 1 or 2 substituents **Y** wherein each **Y** is as defined below;

or

**R<sup>4</sup> and R<sup>5</sup>** may optionally form a heterocyclic ring;

Y is each and independently selected from any of hydrogen, CH<sub>3</sub>; NO<sub>2</sub>;

—(CH<sub>2</sub>)<sub>p1</sub>CF<sub>3</sub>; halogen; —CONR<sup>a</sup>R<sup>b</sup>; —COOR<sup>a</sup>; —COR<sup>a</sup>; —(CH<sub>2</sub>)<sub>p2</sub>NR<sup>a</sup>R<sup>b</sup>;

—(CH<sub>2</sub>)<sub>p3</sub>CH<sub>3</sub>(CH<sub>2</sub>)<sub>p4</sub>SOR<sup>a</sup>R<sup>b</sup>; —(CH<sub>2</sub>)<sub>p5</sub>SO<sub>2</sub>R<sup>a</sup>; —(CH<sub>2</sub>)<sub>p6</sub>SO<sub>2</sub>NR<sup>a</sup>; -O-Aryl;

C<sub>4</sub>-C<sub>8</sub>(alkyl-cycloalkyl) wherein alkyl is C<sub>1</sub>-C<sub>2</sub> alkyl and cycloalkyl is C<sub>3</sub>-C<sub>6</sub>

cycloalkyl; and 1 or 2 heteroaryl(s) having from 5 to 10 atoms and the

heteroatom(s) being selected from any of S, N and O; and wherein

R<sup>a</sup> and R<sup>b</sup> is each and independently selected from hydrogen, a branched or

straight C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkenyl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl; and wherein

p<sup>1</sup>, p<sup>2</sup>, p<sup>3</sup>, p<sup>4</sup>, p<sup>5</sup> and p<sup>6</sup> is each and independently 0, 1 or 2.

Within the scope of the invention are also pharmaceutically acceptable salts of the compounds of the formula (I), as well as isomers, hydrates, isoforms and prodrugs thereof.

Examples of heterocyclic ring systems which may be formed by R<sup>2</sup> and R<sup>3</sup> together include but are not limited to azeridine, pyrrolidine, piperidine, azepine, azocine, their hydrogenated or dehydrogenated derivatives, their aminoderivatives and other aza-heterocycle moieties and their derivatives, such as dihydroimidazoles, di-, tetra- and hexahydropyrimidines and the like.

By "halogen" we mean chloro, fluoro, bromo and iodo.

By "aryl" we mean an aromatic ring having 6 or 10 carbon atoms, such as phenyl and  
5 naphtyl.

By "heteroaryl" we mean an aromatic ring in which one or more of the from 5-10 atoms in  
the ring are elements other than carbon, such as N, S and O.

10 By "isomers" we mean compounds of the formula (I), which differ by the position of their  
functional group and/or orientation. By "orientation" we mean stereoisomers,  
diastereoisomers, regioisomers and enantiomers.

By "isoforms" we mean compounds of the formula (I) which differ by their crystal lattice,  
15 such as crystalline compound and amorphous compounds.

By "prodrug" we mean pharmacologically acceptable derivatives, e.g. esters and amides,  
such that the resulting biotransformation product of the derivative is the active drug. The  
reference by Goodman and Gilman's, *The Pharmacological basis of Therapeutics*, 8th ed.,  
20 McGraw-Hill, Int. Ed. 1992, "Biotransformation of Drugs, p. 13-15, describing prodrugs  
generally, is hereby incorporated.

The novel compounds of the present invention are useful in therapy, especially for the  
treatment of various pain conditions such as chronic pain, acute pain, cancer pain, pain  
25 caused by rheumatoid arthritis, migraine, visceral pain etc. This list should however not be  
interpreted as exhaustive.

Compounds of the invention are useful as immunomodulators, especially for autoimmune diseases, such as arthritis, for skin grafts, organ transplants and similar surgical needs, for collagen diseases, various allergies, for use as anti-tumour agents and anti viral agents.

5 Compounds of the invention are useful in disease states where degeneration or dysfunction of opioid receptors is present or implicated in that paradigm. This may involve the use of isotopically labelled versions of the compounds of the invention in diagnostic techniques and imaging applications such as positron emission tomography (PET).

10 Compounds of the invention are useful for the treatment of diarrhoea, depression, urinary incontinence, various mental illnesses, cough, lung oedema, various gastro-intestinal disorders, spinal injury and drug addiction, including the treatment of alcohol, nicotine, opioid and other drug abuse and for disorders of the sympathetic nervous system for example hypertension.

15 Compounds of the invention are useful as an analgesic agent for use during general anaesthesia and monitored anaesthesia care. Combinations of agents with different properties are often used to achieve a balance of effects needed to maintain the anaesthetic state (eg. Amnesia, analgesia, muscle relaxation and sedation). Included in this combination are inhaled anaesthetics, hypnotics, anxiolytics, neuromuscular blockers and  
20 opioids.

The compounds of the present invention in isotopically labelled form are useful as a diagnostic agent.

25 Also within the scope of the invention is the use of any of the compounds according to the formula (I) above, for the manufacture of a medicament for the treatment of any of the conditions discussed above.

A further aspect of the invention is a method for the treatment of a subject suffering from any of the conditions discussed above, whereby an effective amount of a compound according to the formula (I) above, is administered to a patient in need of such treatment.


5

### Methods of preparation

The compounds of the present invention may be prepared as described in Scheme 1 below.

### 10 General procedure for the preparation of 1,4 or 1,3-guanidinomethyl aminomethyl xylene

Scheme 1



As shown in Scheme 1 above, compounds of the **formula VI** may be obtained from commercially available bis-amino xylylene (**compound 1**).

**Compound 1** is converted into mono-(diBoc)-guanidinomethyl derivative **2** using a protected guanylation reagent such as 1-H-pyrazole-1-(N,N-bis (tert-butoxycarbonyl) carboxamidine in an organic solvent such as THF.

The secondary amine of the **formula III** may be generated using a reductive amination step, where **compound 2** is reacted with an aldehyde **II** in the presence of an acid such as acetic acid or a Lewis acid such as  $ZnCl_2$ , in a protic solvent such as methanol or ethanol in the presence of a reducing agent such as sodium cyanoborohydride.

Compounds of the **formula V** may be obtained by performing an urea formation using **compound III** with a chloroformate of the **formula IV** in a solvent such as methylene chloride and in the presence of a tertiary amine as base, such as triethylamine.

Finally, a compound of the **formula VI** may be obtained by cleavage of the Boc protecting group with an acid such as aqueous hydrochloric acid or by using organic acid such as trifluoroacetic acid in a solvent such a methylene chloride.

The invention will now be described in more detail by way of the following Examples, which are not to be construed as limiting the invention in any way.

#### Step 1 (a)

#### Preparation of 1-(diBoc)-guanidinomethyl-4-aminomethyl benzene (compound 2)

##### Part A

1-H-pyrazole-1-carboxamidine was prepared according to Bernatowicz et.al., J. Org. Chem. 1992, 57, pp.2497-2502, and protected with di-*tert*-butyl

dicarbonate to give 1-H-pyrazole-1-*N,N*-bis(tert-butoxycarbonyl)carboxamide (**compound 1**) according to *Drake et.al, Synth. 1994. pp.579-582.*

5 **Part B**

To a solution of *p*-xylylenediamine (**compound 1**) (30.8 g, 0.226 mol) in THF (300 mL) was added a solution of 1-H-Pyrazole-1-(*N,N*-bis(tert-butoxycarbonyl)carboxamide (35.0 g, 0.113 mol) in THF (100 mL). The solution was stirred at room temperature for 3 h. The solvent was removed under reduced pressure. Water was added to the residue and the  
10 aqueous mixture was extracted with ethyl acetate. The organic layer was washed with brine, dried over MgSO<sub>4</sub> and concentrated. The product (**compound 2**) was purified by column chromatography on silica gel using a mixture of methylene chloride:methanol as the eluent to give 24.3 g (57 % yield) of 1-(diBoc)-guanidinomethyl-4-aminomethyl benzene (**compound 2 where NH<sub>2</sub> is in 4-position**).

15

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.5 (broad s, 1H), 7.32 (s, 4H), 4.65 (d, 2H), 3.89 (s, 2H), 1.5 (s, 9H), 1.48 (s, 9H).

20 **Step 1 (b)**

**Preparation of 1-(diBoc)-guanidinomethyl-3-aminomethyl benzene**

1-(diBoc)-guanidinomethyl-3-aminomethyl benzene was prepared in a similar fashion from *m*-xylylenediamine and of 1-H-Pyrazole-1-(*N,N*-bis(tert-butoxycarbonyl)carboxamide.

25 <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.52 (broad s, 1H), 7.28-7.08 (m, 4H), 4.56 (d, 2H), 3.81 (s, 2H), 1.42 (s, 9H), 1.39 (s, 9H).

**Step 2****Reductive amination: Preparation of 1-(diBoc)-guanidinomethyl-4-[N-(cyclohexylmethyl)]benzene (compound 2 where NH<sub>2</sub> is in 3-position)**

- 5 To a methanolic solution (15 ml) of compound 2 where the amino group is in 4-position) (341 mg, 0.90 mmol) and cyclohexanecarboxaldehyde (111.17 mg, 0.99 mmol) was added zinc chloride (122.79 mg, 0.90 mmol) and sodium cyanoborohydride (67.93 mg, 1.08 mmol). The mixture was stirred over night under nitrogen, wherafter the mixture was diluted with saturated aqueous sodium bicarbonate, and extracted with methylene chloride.
- 10 The organic phase was washed with brine, dried over MgSO<sub>4</sub> and concentrated. This crude product was further purified by silica gel chromatography using CH<sub>2</sub>Cl<sub>2</sub>/MeOH (95:5) as the solvent, to give 164 mg of the pure desired product (compound 2 where NH<sub>2</sub> is in 3-position).

15 **1H NMR (CDCl<sub>3</sub>)**


**δ (ppm):** 0.83 (2H, m, cyclohexane ring); 1.10 (3 H, m, cyclohexane ring); 1.42 (9H, s, boc), 1.46 (9H, s, boc), 1.65 (6H, m, cyclohexane ring), 2.41 (2H, d, *J*=6.8 Hz, C<sub>6</sub>H<sub>11</sub> - CH<sub>2</sub>), 3.72 (2H, s, C<sub>6</sub> H<sub>4</sub> - CH<sub>2</sub>), 4.54 (2H, d, *J*=5.6 Hz, NNCNH - CH<sub>2</sub> - C<sub>6</sub>H<sub>4</sub>), 7.18~7.25 (4H, m, Ar), 8.50 (1H, br, NH - CNN) ppm.

20

Specific examples illustrating the preparation of secondary amines, i.e. intermediates of the formula III, are provided in Table 1 below.



**Table 1**

| Intermediate compound no. | Intermediate compound of the formula III and chemical name                           | Aldehyde [R <sup>3</sup> CHO] | Characterization data  |
|---------------------------|--|-------------------------------|--|
| 3                         |  | cyclohexane carboxaldehyde    | <sup>1</sup> H NMR (CDCl <sub>3</sub> )<br>$\delta$ (ppm): 0.83 (2H, m, cyclohexane ring);<br>1.42 (9H, s, boc),<br>1.46 (9H, s, boc),<br>1.65 (6H, m, cyclohexane ring), 2.41 (2H, d, $J=6.8$ Hz,<br>C <sub>6</sub> H <sub>11</sub> - CH <sub>2</sub> ), 3.72 (2H, s,<br>C <sub>6</sub> H <sub>4</sub> - CH <sub>2</sub> ), 4.54 (2H, d,<br>$J=5.6$ Hz, NNCNH - CH <sub>2</sub> -<br>C <sub>6</sub> H <sub>4</sub> ), 7.18 ~ 7.25 (4H, m, Ar), 8.50 (1H, br, NH - CNN) ppm. |

**Table I (contd.)**




| Intermediate compound no. | Intermediate compound of the formula III and chemical name   | Aldehyde [R <sup>3</sup> CHO] | Characterization data   |
|---------------------------|--|-------------------------------|---|
| 4                         |  <p>1-(diBoc)-guanidinomethyl-4-[N-(2,2-diphenylethyl)aminomethyl] benzene</p> | Diphenylbenzaldehyde          | <p><sup>1</sup>H NMR (CDCl<sub>3</sub>)</p> <p>δ 8.5 (broad s, 1H),</p> <p>7.23-7.14 (m, 14H),</p> <p>4.58 (d, 2H),</p> <p>4.18 (t, 1H), 3.78 (s, 2H),</p> <p>3.22 (d, 2H), 1.50 (s, 9H),</p> <p>1.47 (s, 9H).</p> <p>MS(FAB+): 559 (M+H), 359.</p> |

Table 1 (Contd.)


| Intermediate compound no. | Intermediate compound of the formula III and chemical name  | Aldehyde [R <sup>3</sup> CHO] | Characterization data  |
|---------------------------|---|-------------------------------|--|
| 5                         |  <p>1-(diBoc)-guanidinomethyl-4-[N-(4-chlorobenzyl)aminomethyl] benzene</p> | 4-chlorobenzaldehyde          | <sup>1</sup> H NMR (DMSO-d <sub>6</sub> )<br>δ 8.6 (broad t, 1H),<br>7.42-7.1 (m, 10H),<br>4.52 (t, 2H), 3.7 (s, 2H),<br>3.2 (s, 2H), 1.5 (s, 9H),<br>1.45 (s, 9H).<br><br>MS(FAB+): 503 (M+H),<br>403, 303. |

73 300


Table 1 (contd.)

| Intermediate compound no. | Intermediate compound of the formula III and chemical name  | Aldehyde [R <sup>3</sup> CHO] | Characterization data   |
|---------------------------|---|-------------------------------|---|
| 6                         |  <p>1-(diBoc)-guanidinomethyl-4-[N-(benzyl)aminomethyl] benzene</p> | Benzaldehyde                  | <sup>1</sup> H NMR (CDCl <sub>3</sub> )<br>δ 7.2 (m, 9H), 4.5 (s, 2H),<br>4.15 (t, 2H), 3.7 (s, 2H),<br>1.5 (s, 18H). |

**Table 1 (Contd.)**

| Intermediate compound no. | Intermediate compound of the formula III and chemical name  | Aldehyde [R <sup>3</sup> CHO] | Characterization data   |
|---------------------------|---|-------------------------------|---|
| 7                         |  <p>1-(diBoc)-guanidinomethyl-4-[N-(2-chlorobenzyl)aminomethyl] benzene</p> | 2-chlorobenzaldehyde          | <sup>1</sup> H NMR (CDCl <sub>3</sub> )<br>δ 8.62 (broad s, 1H),<br>7.48-7.1 (m, 8H),<br>4.62 (d, 2H), 3.95 (s, 2H),<br>3.85 (s, 2H), 1.53 (s, 9H),<br>1.5 (s, 9H). |

**Table 1 (contd.)**

| Intermediate compound no. | Intermediate compound of the formula III and chemical name  | Aldehyde [ $R^3CHO$ ] | Characterization data  |
|---------------------------|---|-----------------------|--|
| 8                         |  <p>1-(diBoc)-guanidinomethyl-4-[N-(3-chlorobenzyl)aminomethyl] benzene</p> | 3-chlorobenzaldehyde  | $^1H$ NMR ( $CDCl_3$ )<br>$\delta$ 8.56 (broad t, 1H),<br>7.4-7.15 (m, 8H),<br>4.6 (d, 2H), 3.8 (s, 4H),<br>1.56 (s, 9H), 1.52 (s, 9H).<br><br>MS(APCI): 503 (M+H),<br>403, 303. |

**Table 1 (Contd.)**




| Intermediate compound no. | Intermediate compound of the formula III and chemical name   | Aldehyde [R <sup>3</sup> CHO] | Characterization data         |
|---------------------------|--|-------------------------------|-------------------------------|
| 9                         |  <p>1-(diBoc)-guanidinomethyl-3-[N-(2,2-diphenylethyl)aminomethyl] benzene</p> | Diphenylacetaldehyde          | MS(ES+): 559 (M+H), 459, 359. |

Table 1 (contd.)

| Intermediate compound no. | Intermediate compound of the formula III and chemical name  | Aldehyde [R <sup>3</sup> CHO] | Characterization data  |
|---------------------------|---|-------------------------------|--|
| 10                        |  <p>1-(diBoc)-guanidinomethyl-3-[N-(4-chlorobenzyl)aminomethyl] benzene</p> | 4-chlorobenzaldehyde          | <sup>1</sup> H NMR (CDCl <sub>3</sub> )<br>δ 8.48 (broad s, 1H),<br>7.3-6.96 (m, 8H),<br>4.68-4.32 (m, 4H),<br>3.78-3.6 (m, 2H). |



**Schem 2**



Alternatively, as shown in Scheme 2, compounds of the formula (XI) may be obtained by using compounds of the formula (VIII) wherein  $X=CN$  and  $Y=CHO$ , as a starting material.

A reductive amination using a primary amine with compound (VIII) in the presence of an acid such as acetic acid, and in the presence of a reducing agent such as sodium cyanoborohydride in a solvent such as methanol or ethanol, provides a compound of the formula (IX).

Compounds of the formula (X) may be obtained by performing an urea reaction using compounds of the formula (IX) with a chloroformate of the formula (V) in a solvent such as methylene chloride and in the presence of a tertiary amine as base, such as triethylamine.

Compounds of the formula (XI) may be prepared by a reduction of the nitrile function in formula (X), using a reduction agent such as borane-THF complex in a solvent such as THF.

Compounds of the formula (XV) may be prepared by reacting compounds of the formula (VIII) wherein  $X=CH_2Br$  and  $Y=CN$ , with an amine in a solvent such as acetonitrile, providing a compound of the formula (XII). A reduction of the nitrile function using a reducing agent such as borane-THF complex in a solvent such as THF, provides the primary amine of the formula (XIII).

A reductive amination step of (XIII) as described above, provides a compound of the formula (XIV). Finally, urea formation of the secondary amine (XIV) as described above, provides a compound of the formula (XV).

Alternatively, compounds of the formula (XV) may be prepared by using a monoprotected dialdehyde such as a compound of the formula (VIII) wherein  $X=CH(OEt)_2$  and  $Y=CHO$ , and a reductive amination in the presence of a reducing agent such as sodium cyanoborohydride in a solvent such as methanol or ethanol. Urea formation as described

above provides a compound of the formula (XVII). Hydrolysis of the diethyl acetal function in compound (XVII) using an acid such as TFA in a solvent such as methylene chloride, provides the corresponding aldehyde (XVIII).


- 5 Finally, a reductive amination as described before, provides compounds of the formula (XV).

The invention will now be described in more detail by the following Examples, which are  
10 not to be construed as limiting the invention.

**EXAMPLE 1**

**Preparation of 1-N-(cyclohexylmethyl)-N-(N-methyl-N-phenylcarbamoyl)-aminomethyl-4-guanidinomethyl-benzene (Compound 12)**

Compound 12 of the present Example was prepared by following the synthetic route described in Scheme 2 below.



- 10 To a solution of compound 3 (164 mg, 0.35 mmol) in methylene chloride (10 ml) was added N-methyl-N-phenyl carbamoylchloride (120.78 mg, 0.71 mmol) and triethylamine (71.90 mg, 0.71 mmol). The mixture was stirred at room temperature for 3 h, washed with a saturated  $\text{NH}_4\text{Cl}$  aqueous solution and brine, dried over  $\text{MgSO}_4$  and concentrated to give

the crude product (**compound 11**). This crude compound was used directly without purification for the preparation of compound 12. It was dissolved in dry methylene chloride (3 ml), 1.5 ml of TFA was added and the reaction mixture was stirred at room temperature for 1 hour. The excess of solvent and TFA was evaporated, the residue was purified by reverse phase preparative HPLC to give the pure desired product (100 mg, 71 % in 2 steps).

$^1\text{H NMR}$  ( $\text{CDCl}_3$ )  $\delta$  (ppm): 0.70 (2H, m, cyclohexane ring); 1.10 (3H, m, cyclohexane ring); 1.41 (3H, m, cyclohexane ring); 1.53 (3H, m, cyclohexane ring); 2.72 (2H, d,  $J=6.4$  Hz,  $\text{C}_6\text{H}_7 - \text{CH}_2$ ), 3.00 (3H, s,  $\text{N-CH}_3$ ), 4.13 (2H, s,  $\text{C}_6\text{H}_4 - \text{CH}_2$ ), 4.24 (2H, d,  $J=4.8$  Hz,  $\text{NH-CH}_2\text{-Ph}$ ), 6.93~7.25 (9H, m, Ar), 8.13 (1H, br,  $\text{NH-C=N}$ ).

**MS Observed (CI):** 408.45 (MH+).

The following specific compounds were prepared by following the synthesis description described above.

**EXAMPLES 2-7**

The following compounds were prepared by using the same procedure as described in Example 1, but using the intermediate and acid chloride indicated in Table 2 below.

Table 2


| Example | Structure and chemical name   | Intermediate | Carbamoyl chloride or Isocyanate    | Physical Characterization  |
|---------|---|--------------|-------------------------------------|--|
| 2       |  <p style="text-align: center;">(13)</p> <p>1-N-[(4-chlorobenzyl)-N-(N-methyl-N-phenylcarbamoyl)]-aminomethyl-4-guanidino-benzene</p> | 5            | N-methyl-N-phenylcarbamoyl chloride | <sup>1</sup> H NMR (DMSO-d <sub>6</sub> ) δ 8.0 (t, 1H), 8.6-8.2 (m, 14 H), 4.4 (d, 2H), 4.2 (d, 4H), 3.1 (s, 3H). |

Table 2 (contd.)


| Example | Structure and chemical name  | Intermediate | Carbamoyl chloride         | Physical Characterization  |
|---------|--|--------------|----------------------------|--|
| 3       |  <p style="text-align: center;">(14)</p> <p>1-N-[(4-chlorobenzyl)-N-(N,N-dimethylcarbamoyl)]-N-(N,N-dimethylcarbamoyl)-4-guanidinomethyl-benzene</p> | 5            | Dimethylcarbamoyl-chloride | <sup>1</sup> H NMR (DMSO-d <sub>6</sub> )<br>δ 8.1 (broad, 1H), 7.4 (d, 2H), 7.25 (d, 2H), 7.15 (m, 4H), 4.35 (d, 2H), 4.15 (d, 4H), 2.85 (s, 6H).<br><br>MS(APCI): 374 (M+H). |

Table 2contd.



| Example | Structure and chemical name   | Interme-<br>diate | Carbamoyl chloride                  | Physical Characterization  |
|---------|---|-------------------|-------------------------------------|--|
| 4       |  <p data-bbox="998 1459 1031 1522">(15)</p> <p data-bbox="1063 1291 1218 1764">1-N-[(4-chlorobenzyl)-N-(N-methyl-N-phenylcarbamoyl)]-aminomethyl-3-guanidinomethyl-benzene</p> | 10                | N-methyl-N-phenylcarbamoyl chloride | <p data-bbox="576 294 730 756"><sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.05-7.5 (m, 13H), 4.4 (broad s, 2H), 4.15 (s, 2H), 3.35 (s, 3H), 3.05 (d, 2H).</p> <p data-bbox="803 472 836 756">MS(APCI): 436 (M+H)</p> |



Table 2 (contd.)

| Example | Structure and chemical name  | Intermedate | Carbamoyl chloride      | Physical Characterization   |
|---------|--|-------------|-------------------------|---|
| 5       |  <p style="text-align: center;">(16)</p> <p>1-N-[(4-chlorobenzyl)-N-(3-nitrophenylcarbamoyl)]-N-(3-guanidinomethyl)-benzene</p> | 5           | 3-nitrophenylisocyanate | <p><sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 9.2 (s, 1H), 8.6 (s, 1H), 8.1 (broad, 1H), 8.0 (d, 1H), 7.7 (d, 1H), 7.55 (t, 1H), 7.4 (d, 1H), 7.2 (m, 6H), 4.7 (d, 4H), 4.3 (d, 2H).</p> <p>MS(APCI): 466 (M+H)</p> |

**Table 2 (contd.)**


| Example | Structure and chemical name  | Interme-<br>diate | Carbamoyl chloride         | Physical Characterization  |
|---------|--|-------------------|----------------------------|--|
| 6       |  <p style="text-align: right;">(17)</p> <p>1-N-[(4-chlorobenzyl)-N-(4-phenoxyphenyl)-aminomethyl]-4-phenoxycarbonyl]-aminomethyl-4-guanidinomethyl-benzene</p> | 5                 | 4-phenoxyphenyl-isocyanate | <sup>1</sup> H NMR (DMSO-d <sub>6</sub> ) δ 8.8 (s, 1H),<br>7.8-7.5 (m, 12H), 7.1 (t, 1H),<br>6.9 (d, 4H), 4.5 (d, 4H), 4.3 (d, 2H).<br><br>MS(APCI): 514. (M+H) |

Table 2 (contd.)



| Example | Structure and chemical name   | Intermediate | Carbamoyl chloride                  | Physical Characterization   |
|---------|---|--------------|-------------------------------------|---|
| 7       |  <p>(18)</p> <p>1-N-[(2-chlorobenzyl)-N-(N-methyl-N-phenylcarbamoyl)]-aminomethyl-4-guanidinomethyl-benzene</p> | 7            | N-methyl-N-phenylcarbamoyl chloride | <sup>1</sup> H NMR (DMSO-d <sub>6</sub> ) δ 8.2 (broad, 1H), 7.4-6.95 (m, 13H), 4.5 (s, 2H), 4.2 (d, 4H), 3.1 (s, 3H).<br>MS(APCI): 436. (M+H). |

Table 2 (contd.)


| Example | Structure and chemical name  | Intermediate | Carbamoyl chloride                  | Physical Characterization  |
|---------|--|--------------|-------------------------------------|--|
| 8       | <p>(19)</p> <p>1-N-[(3-chlorobenzyl)-N-(N-methyl-N-phenylcarbamoyl)]-aminomethyl-4-guanidinomethyl-benzene</p> | 8            | N-methyl-N-phenylcarbamoyl chloride | <sup>1</sup> H NMR (DMSO-d <sub>6</sub> )<br>δ 8.2 (broad, 1H), 7.4-7.1 (m, 13H),<br>4.35 (d, 2H), 4.15 (s, 4H), 3.1 (s, 3H).<br><br>MS(APCI): 436. (M+H). |

Table 2 (contd.)

| Example | Structure and chemical name   | Intermediate | Carbamoyl chloride                   | Physical Characterization  |
|---------|---|--------------|--------------------------------------|--|
| 9       |  <p>(20)</p> <p>1-N-[(benzyl)-N-(N-methyl-N-phenylcarbamoyl)]-aminomethyl-4-guanidinomethyl-benzene</p> | 6            | N-methyl-N-phenyl-carbamoyl chloride | <sup>1</sup> H NMR (DMSO-d <sub>6</sub> ) δ 8.1(t, 1H), 7.3 (m, 8H), 7.05 (m, 6H), 4.45 (d, 2H), 4.05 (d, 4H), 3.15 (s, 3H). |

**EXAMPLE 10**

**Preparation of 1-N-[(2,4-dichlorobenzyl)-N-(N-methyl-N-phenylcarbamoyl)]-aminomethyl-4-guanidinomethyl-benzene (Compound 21)**

**(21)**

5

Following the same procedure as described in Example 1, step 2 but substituting 2,4-dichlorobenzaldehyde for cyclohexanecarboxaldehyde, the title **compound 21** was  
10 obtained.

$^1\text{H NMR}$  (DMSO- $d_6$ )  $\delta$  8.2 (broad, 1H), 7.5-7.1 (m, 12H), 4.5 (d, 2H), 4.2 (s, 4H), 3.1 (s, 3H).


MS(APCI): 469.95 (M+H).

15

**EXAMPLE 11****Preparation of 1-N-[(4-chlorobenzyl)-N-(N-methyl-N-phenylcarbamoyl)]-aminomethyl-4-aminomethyl-benzene (Compound 25)**

5

Compound 25 of the present Example was prepared by following the procedure described in Scheme 3 below.



10

**Step 1****Preparation of N-(4-chlorobenzyl)-4-cyanobenzyl amine (compound 22)**

To a methanolic solution (20 mL) of 4-chlorobenzylamine (1.02 g, 7.2 mmol) was added successively ZnCl<sub>2</sub> (.981 g, 7.2 mmol), 4-cyanobenzaldehyde (1.007 g, 7.3 mmol), and NaCNBH<sub>3</sub> (0.452 g, 7.2 mmol). The reaction mixture was stirred at r.t. for 2 days. It was diluted with aq. sodium bicarbonate and the reaction mixture was extracted with methylene chloride. The combined organic extracts were washed with brine, dried over magnesium sulfate and concentrated. The product (**compound 22**) was purified by silica gel chromatography: 1.43 g (77%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.45 (d, 2H), 7.28 (d, 2H), 7.10 (s, 4H), 3.68 (s, 2H), 3.58 (s, 2H).

**Step 2****Preparation of 1-N-[(4-chlorobenzyl)-N-(N-methyl-N-phenylcarbamoyl)]-aminomethyl-4-cyanobenzene (compound 24)**

To a solution of N-(4-chlorobenzyl)-4-cyanobenzyl amine (**compound 22**) (1.43 g, 5.57 mmol) in dioxane (20 mL) was added N-methyl-N-phenyl carbamoyl chloride (**compound 23**) (1.039 g, 6.12 mmol) and triethylamine (0.853 mL, 6.12 mmol). The reaction mixture was stirred at r.t. for 1 day, then it was diluted with ethyl acetate and washed with 10% HCl, saturated sodium bicarbonate, water, brine, dried over MgSO<sub>4</sub> and concentrated to give **compound 24**: 1.95 g (89%).

**Step 3****Preparation of 1-N-[(4-chlorobenzyl)-N-(N-methyl-N-phenylcarbamoyl)]-aminomethyl-4-aminomethyl-benzene (compound 25)**

To a THF (6 mL) solution of 1-N-[(4-chlorobenzyl)-N',N'-(methyl, phenyl)carbamoyl]-aminomethyl-4-cyanobenzene (**compound 24**) (0.39 g, 1 mmol) was added a 1M BH<sub>3</sub>.THF complex (2.2 mL). The mixture was heated at 90 °C overnight. Then a 2.55 M HCl in methanol (3 mL) was added and the reaction mixture was heated at reflux for 1 h. The mixture was diluted with water and extracted with ethyl acetate. The organic layer




was dried over  $MgSO_4$  and concentrated. The product (**compound 25**) was purified by preparative TLC using methanol/methylene chloride/ammonium hydroxide as the eluent.

$^1H$  NMR ( $CDCl_3$ )  $\delta$  7.20-6.84 (m, 13H), 4.00 (s, 4H), 3.70 (s, 2H), 3.05 (s, 3H), 2.05 (broad s, 2H). MS: 394. (M+H).

### EXAMPLE 12

#### Preparation of 1-N-[4-chlorobenzyl]-N-(N-methyl-N-phenylcarbamoyl)-aminomethyl-4-(N-pyrrolidinomethyl)-benzene (compound 32)

The compound 32 of Example 11 was prepared by following the procedure described in Scheme 4 below.



**Step 1****Preparation of 4-cyano-1-N-pyrrolinomethyl benzene (compound 28)**

4-cyano benzyl bromide (**compound 26**) (20.0 g, 0.102 mol) was dissolved in  
5 acetonitrile (100 mL) and added to a cooled ( $0^{\circ}\text{C}$ ) solution of pyrrolidine (**compound 27**)  
(8.5 g, 0.12 mol) in acetonitrile. The mixture was stirred at RT for 3 days. The reaction  
mixture was acidified with 4N HCl and washed with ethyl acetate. The aqueous layer was  
basified with 20% sodium hydroxide and extracted with ethyl acetate. The organic layer  
was concentrated to give product (**compound 28**) (11.0 g, 58%) which was carried on to  
10 the next step with no further purification.

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 7.6 (d, 2H), 8.4 (d, 2H), 3.55 (s, 2H), 2.55 (t, 2H), 1.7 (t, 2H). MS:  
187. (M+H)

**Step 2****Preparation of 4-N-aminomethyl-1-N-pyrrolinomethyl benzene (compound 29)**

4-cyanomethyl-1-N-pyrrolinomethyl benzene (**compound 28**) (11 g, 59 mmol) was  
dissolved in dry THF (20 mL). To this solution was added a 1M solution of borane/THF  
complex (180 mL). The mixture was refluxed overnight. The solution was then cooled to  
r.t. and a solution of 3N HCl in methanol (120 mL) was added dropwise. The mixture was  
20 again refluxed overnight. After cooling to room temperature, the product (**compound 29**)  
fell out of solution as white precipitate and was collected and washed with THF: 11.6 g.

$^1\text{H NMR}$  ( $\text{CDCl}_3$ )  $\delta$  8.8 (broad, 1H), 7.6 (d, 2H), 7.4 (d, 2H), 4.4 (s, 2H), 4.0 (s, 2H), 3.3  
(t, 2H), 3.0 (t, 2H), 2.0 (t, 4H).

25 MS: 191. (M+H)

**Step 3****Preparation of N-(4-chlorobenzyl)-1-(1-N-pyrrolidinomethyl)-benzylamine (compound 31)**

5 Following the same procedure as described for Example 10, step 1, but substituting 4-chlorobenzylamine for compound 29 and 4-cyanobenzaldehyde for 4-chlorobenzaldehyde, the title compound was obtained.

MS: 357 (M+H)

10

**Step 4****Preparation of 1-N-[(4-chlorobenzyl)-N-(N-methyl-N-phenylcarbamoyl)]-aminomethyl-4-(1-N-pyrrolidinomethyl)-benzene (compound 32)**

15

Following the procedure described in Example 10, step 2, but substituting compound 22 for compound 31, the title compound was obtained.


20 <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.55 (d, 2H), 7.45 (d, 2H), 7.4 (d, 2H), 7.2 (d, 2H), 7.0 (d, 2H), 6.95 (d, 2H), 4.3 (d, 4H), 4.1 (s, 2H), 3.25 (s, 3H), 3.1 (t, 4H), 2.05 (t, 4H).

MS 448. (M+H)

25

**EXAMPLE 13****Preparation of 1-N-[(4-chlorobenzyl)-N-(N-methyl-N-phenylcarbamoyl)]-aminomethyl-4-(N,N-dimethyl)-aminomethyl-benzene (compound 35)**

The compound 35 of Example 12 was prepared by following the procedure described in  
30 Scheme 5 below.

**Step 1****Preparation of 1-N-(4-chlorobenzyl)-4-(N,N-dimethyl)aminomethyl benzyl amine****5 (compound 34)**

A round bottom flask was charged with 1-N-aminomethyl-4-N,N-dimethylaminomethyl benzene hydrochloride (**compound 33**) (2.36 g, 10 mmol), 4-chlorobenzaldehyde (**compound 30**) (1.51 g, 11 mmol, 97% pure), and methanol (40 mL). The mixture was stirred at r.t. for 20 minutes, then solid NaCNBH<sub>3</sub> was added and the reaction mixture was stirred at r.t. overnight. It was diluted with aqueous sodium bicarbonate, and extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over MgSO<sub>4</sub> and concentrated to give 1.29 g of (**compound 34**) as an oily residue.

**Step 2****15 Preparation of 1-N-[(4-chlorobenzyl)-N-(N-methyl-N-phenylcarbamoyl)]-aminomethyl-4-(N,N-dimethyl)aminomethyl-benzene (compound 35)**

To a solution of 1-N-(4-chlorobenzyl)-4-(N,N-dimethyl)aminomethyl benzyl amine (**compound 34**) (0.562 g, 1.95 mmol) in dioxane (8 mL) was added triethylamine (0.326 mL, 2.34 mmol) and N-methyl,N-phenyl carbamoyl chloride (**compound 23**) (0.395 g, 2.34 mmol). The mixture was stirred at r.t. overnight, then diluted with aqueous sodium bicarbonate and extracted with ethyl acetate. The organic extracts were washed with brine, dried over MgSO<sub>4</sub> and concentrated to a colorless oil. The product (**compound 35**) was

purified by silica gel chromatography using methanol/ethyl acetate/ammonium hydroxide as the eluent.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.40-6.95 (m, 13H), 4.20 (s, 4H), 3.42 (s, 2H), 3.25 (s, 3H), 2.32 (s, 6H).

MS: 422. (M+H).


#### **EXAMPLE 14**

10 **Preparation of 1-N-[4-chlorobenzyl)-N-(N-methyl-N-phenylcarbamoyl)]-aminomethyl-4-(N-methyl)-aminomethyl-benzene (Compound 40)**

The compound 40 of Example 13 was prepared by following the procedure described in Scheme 6 below.

15

Scheme 6

**Step 1****Preparation of 1-diethyl acetal-4-N-(4-chlorobenzyl) benzyl amine (compound 37)**

- 5 To a solution of *tere*-phthalaldehyde *mono*-(diethyl acetal) (compound 36) (0.416 g, 2 mmol) and 4-chlorobenzylamine (compound 21) (0.283 g, 2 mmol) in methanol (5 mL, containing 1% glacial acetic acid v/v) was added solid  $\text{NaCNBH}_3$ . The reaction mixture was stirred at r.t. overnight. It was diluted with aqueous sodium bicarbonate and extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over
- 10  $\text{MgSO}_4$  and concentrated to an oil, (compound 37): .61 g (91%). MS: 334.02 (M+H).

## Step 2

### Preparation of 1-N-[(4-chlorobenzyl)-N-(N-methyl-N-phenylcarbamoyl)]-aminomethyl-4-carboxaldehyde (compound 39)

5 To a solution of 1-diethyl acetal-4-N-(4-chlorobenzyl) benzyl amine (**compound 37**) (0.60 g, 1.8 mmol) in dioxane (5 mL) was added triethylamine (0.279 mL, 2 mmol) and N-methyl,N-phenyl carbamoyl chloride (**compound 23**) (0.336g, 1.98 mmol). The mixture was stirred at r.t. overnight, then diluted with aqueous sodium bicarbonate and extracted with ethyl acetate. The organic extracts were washed with brine, dried over MgSO<sub>4</sub> and  
10 concentrated to give (**compound 38**) : 0.69 g.

The acetal (**compound 38**) (0.67 g, 1.42 mmol) was dissolved in 50% trifluoroacetic acid/methylene chloride (5 mL) and stirred at r.t. for 3.5 h. The mixture was concentrated to an oily residue which was redissolved in CH<sub>2</sub>Cl<sub>2</sub> and washed with sodium bicarbonate,  
15 brine, dried over MgSO<sub>4</sub> and concentrated to an oil, (**compound 39**): 0.504 g.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 9.80 (s, 1H), 7.72-6.68 (m, 13H), 4.10 (s, 2H), 3.97 (s, 2H), 3.10 (s, 3H).

## 20 Step 3

### Preparation of 1-N-[(4-chlorobenzyl)-N-(N-methyl-N-phenylcarbamoyl)]-aminomethyl-4-(N-methyl)-aminomethyl-benzene (compound 40)

To a methanolic solution (5 mL) of **compound 39** (0.50 g, 1.28 mmol) was added a solution of 2M methylamine in methanol (0.7 mL, 1.40 mmol), and glacial acetic acid  
25 (0.05 mL). Solid NaCNBH<sub>3</sub> (0.08 g, 1.28 mmol) was then added and the mixture was stirred at r.t. overnight. It was diluted with aqueous sodium bicarbonate and extracted with ethyl acetate. The organic extracts were washed with brine, dried and concentrated. The product (**compound 40**) was purified by silica gel chromatography, using a mixture of methanol/methylene chloride/ammonium hydroxide as the eluent.  
30

$^1\text{H NMR (CDCl}_3\text{)}$   $\delta$  7.40-6.95 (m, 13H), 4.19 (s, 2H), 4.16 (s, 2H), 3.82 (s, 2H), 3.19 (s, 3H), 2.46 (s, 3H).

MS: 408.03. (M+H).

5 Pharmaceutical compositions

The novel compounds according to the present invention may be administered orally, intramuscularly, subcutaneously, topically, intranasally, intraperitoneally, intrathoracically, intravenously, epidurally, intrathecally, intracerebroventricularly and by injection into the  
10 joints.

A preferred route of administration is orally, intravenously or intramuscularly.

The dosage will depend on the route of administration, the severity of the disease, age and  
15 weight of the patient and other factors normally considered by the attending physician, when determining the individual regimen and dosage level as the most appropriate for a particular patient.

For preparing pharmaceutical compositions from the compounds of this invention, inert,  
20 pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include powders, tablets, dispersible granules, capsules, cachets, and suppositories.

A solid carrier can be one or more substances which may also act as diluents, flavoring  
agents, solubilizers, lubricants, suspending agents, binders, or tablet disintegrating agents;  
25 it can also be an encapsulating material.

In powders, the carrier is a finely divided solid which is in a mixture with the finely divided  
active component. In tablets, the active component is mixed with the carrier having the  
necessary binding properties in suitable proportions and compacted in the shape and size  
30 desired.



For preparing suppository compositions, a low-melting wax such as a mixture of fatty acid glycerides and cocoa butter is first melted and the active ingredient is dispersed therein by, for example, stirring. The molten homogeneous mixture is then poured into convenient  
5 sized molds and allowed to cool and solidify.

Suitable carriers are magnesium carbonate, magnesium stearate, talc, lactose, sugar, pectin, dextrin, starch, tragacanth, methyl cellulose, sodium carboxymethyl cellulose, a low-melting wax, cocoa butter, and the like.

10

Pharmaceutically acceptable salts are acetate, benzenesulfonate, benzoate, bicarbonate, bitartrate, bromide, calcium acetate, camsylate, carbonate, chloride, citrate, dihydrochloride, edetate, edisylate, estolate, esylate, fumarate, gluaptate, gluconate, glutamate, glycolylarsanilate, hexylresorcinate, hydrabamine, hydrobromide,  
15 hydrochloride, hydroxynaphthoate, iodide, isethionate, lactate, lactobionate, malate, maleate, mandelate mesylate, methylbromide, methylnitrate, methylsulfate, mucate, napsylate, nitrate, pamoate (embonate), pantothenate, phosphate/diphosphate, polygalacturonate, salicylate, stearate, subacetate, succinate, sulfate, tannate, tartrate, teoclate, triethiodide, benzathine, chlorprocaine, choline, diethanolamine,  
20 ethylenediamine, meglumine, procaine, aluminium, calcium, lithium, magnesium, potassium, sodium, and zinc.

Preferred pharmaceutically acceptable salts are the hydrochlorides, trifluoroacetates and bitartrates.

The term composition is intended to include the formulation of the active component with encapsulating material as a carrier providing a capsule in which the active component (with or without other carriers) is surrounded by a carrier which is thus in association with it.

5 Similarly, cachets are included.

Tablets, powders, cachets, and capsules can be used as solid dosage forms suitable for oral administration.

10 Liquid from compositions include solutions, suspensions, and emulsions. Sterile water or water-propylene glycol solutions of the active compounds may be mentioned as an example of liquid preparations suitable for parenteral administration. Liquid compositions can also be formulated in solution in aqueous polyethylene glycol solution.

15 Aqueous solutions for oral administration can be prepared by dissolving the active component in water and adding suitable colorants, flavoring agents, stabilizers, and thickening agents as desired. Aqueous suspensions for oral use can be made by dispersing the finely divided active component in water together with a viscous material such as natural synthetic gums, resins, methyl cellulose, sodium carboxymethyl cellulose, and other  
20 suspending agents known to the pharmaceutical formulation art.

Preferably the pharmaceutical composition is in unit dosage form. In such form, the composition is divided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing  
25 discrete quantities of the preparations, for example, packeted tablets, capsules, and powders in vials or ampoules. The unit dosage form can also be a capsule, cachet, or tablet itself, or it can be the appropriate number of any of these packaged forms.

## BIOLOGICAL EVALUATION

### A) IN VITRO MODEL

#### Cell culture

5

Human 293S cells expressing cloned human  $\mu$ ,  $\delta$ , and  $\kappa$  receptors and neomycin resistance were grown in suspension at 37°C and 5% CO<sub>2</sub> in shaker flasks containing calcium-free DMEM 10% FBS, 5% BCS, 0.1% Pluronic F-68, and 600  $\mu$ g/ml geneticin.

#### 10 Membrane preparation

Cells were pelleted and resuspended in lysis buffer (50 mM Tris, pH 7.0, 2.5 mM EDTA, with PMSF added just prior to use to 0.1 mM from a 0.1 M stock in ethanol), incubated on ice for 15 min, then homogenized with a polytron for 30 sec. The suspension was spun at 15 1000g (max) for 10 min at 4°C. The supernatant was saved on ice and the pellets resuspended and spun as before. The supernatants from both spins were combined and spun at 46,000 g(max) for 30 min. The pellets were resuspended in cold Tris buffer (50 mM Tris/Cl, pH 7.0) and spun again. The final pellets were resuspended in membrane buffer ( 50 mM Tris, 0.32 M sucrose, pH 7.0). Aliquots (1 ml) in polypropylene tubes were 20 frozen in dry ice/ethanol and stored at -70°C until use. The protein concentrations were determined by a modified Lowry assay with SDS.

#### Binding assays

25 Membranes were thawed at 37°C, cooled on ice, passed 3 times through a 25-gauge needle, and diluted into binding buffer (50 mM Tris, 3 mM MgCl<sub>2</sub>, 1 mg/ml BSA (Sigma A-7888), pH 7.4, which was stored at 4°C after filtration through a 0.22  $\mu$ m filter, and to which had been freshly added 5  $\mu$ g/ml aprotinin, 10  $\mu$ M bestatin, 10  $\mu$ M diprotin A, no DTT). Aliquots of 100  $\mu$ l (for  $\mu$ g protein, see Table 1) were added to iced 12x75 mm 30 polypropylene tubes containing 100  $\mu$ l of the appropriate radioligand (see Table 1) and

100  $\mu$ l of test peptides at various concentrations. Total (TB) and nonspecific (NS) binding were determined in the absence and presence of 10  $\mu$ M naloxone respectively. The tubes were vortexed and incubated at 25°C for 60-75 min, after which time the contents are rapidly vacuum-filtered and washed with about 12 ml/tube iced wash buffer (50 mM Tris, pH 7.0, 3 mM  $MgCl_2$ ) through GF/B filters (Whatman) presoaked for at least 2h in 0.1% polyethyleneimine. The radioactivity (dpm) retained on the filters was measured with a beta counter after soaking the filters for at least 12h in minivials containing 6-7 ml scintillation fluid. If the assay is set up in 96-place deep well plates, the filtration is over 96-place PEI-soaked unifilters, which were washed with 3 x 1 ml wash buffer, and dried in an oven at 55°C for 2h. The filter plates were counted in a TopCount (Packard) after adding 50  $\mu$ l MS-20 scintillation fluid/well.

#### Data analysis

15 The specific binding (SB) was calculated as TB-NS, and the SB in the presence of various test peptides was expressed as percentage of control SB. Values of  $IC_{50}$  and Hill coefficient ( $n_H$ ) for ligands in displacing specifically bound radioligand were calculated from logit plots or curve fitting programs such as Ligand, GraphPad Prism, SigmaPlot, or ReceptorFit. Values of  $K_i$  were calculated from the Cheng-Prussoff equation. Mean  $\pm$  20 S.E.M. values of  $IC_{50}$ ,  $K_i$  and  $n_H$  were reported for ligands tested in at least three displacement curves.

### Receptor saturation experiments

Radioligand  $K_d$  values were determined by performing the binding assays on cell membranes with the appropriate radioligands at concentrations ranging from 0.2 to 5 times the estimated  $K_d$  (up to 10 times if amounts of radioligand required are feasible). The specific radioligand binding was expressed as pmole/mg membrane protein. Values of  $K_d$  and  $B_{max}$  from individual experiments were obtained from nonlinear fits of specifically bound (B) vs. nM free (F) radioligand from individual according to a one-site model.

### 10 B) BIOLOGICAL MODEL (IN VIVO MODEL)

FREUND'S COMPLETE ADJUVANT (FCA), AND SCIATIC NERVE CUFF INDUCED MECHANO-ALLODYNIA IN RAT

#### Animals

15 Male Sprague-Dawley rats (Charles River, St-Constant, Canada) weighing 175-200g at the time of surgery were used. They were housed in groups of three in rooms thermostatically maintained at 20° C with a 12:12 hr light/dark cycle, and with free access to food and water. After arrival, the animals were allowed to acclimatize for at least 2 days before surgery. The experiments were approved by the appropriate Medical Ethical Committee  
20 for animal studies.

### EXPERIMENTAL PROCEDURE

#### FREUND'S COMPLETE ADJUVANT

The rats were first anesthetized in a Halothane chamber after which 10µl of FCA was  
25 injected s.c. into the dorsal region of the left foot, between the second and third external digits. The animals were then allowed to recover from anesthesia under observation in their home cage.

### SCIATIC NERVE CUFF

The animals were prepared according to the method described by Mosconi and Kruger (1996). Rats were anesthetized with a mixture of Ketamine / Xylazine i.p. (2ml/kg) and placed on their right side and an incision made over, and along the axis of, the lateral  
5 aspect of the left femur. The muscles of the upper quadriceps were teased apart to reveal the sciatic nerve on which a plastic cuff (PE-60 tubing, 2mm long) was placed around. The wound was then closed in two layers with 3-0 vicryl and silk sutures.

### DETERMINATION OF MECHANO-ALLODYNIA USING VON FREY TESTING

10 Testing was performed between 08:00 and 16:00h using the method described by Chaplan et al. (1994). Rats were placed in Plexiglas cages on top of a wire mesh bottom which allowed access to the paw, and were left to habituate for 10-15 min. The area tested was the mid-plantar left hind paw, avoiding the less sensitive foot pads. The paw was touched with a series of 8 Von Frey hairs with logarithmically incremental stiffness (0.41, 0.69,  
15 1.20, 2.04, 3.63, 5.50, 8.51, and 15.14 grams; Stoelting, III, USA). The von Frey hair was applied from underneath the mesh floor perpendicular to the plantar surface with sufficient force to cause a slight buckling against the paw, and held for approximately 6-8 seconds. A positive response was noted if the paw was sharply withdrawn. Flinching immediately upon removal of the hair was also considered a positive response. Ambulation was  
20 considered an ambiguous response, and in such cases the stimulus was repeated.

### TESTING PROTOCOL

The animals were tested on postoperative day 1 for the FCA-treated group and on post-operative day 7 for the Sciatic Nerve Cuff group. The 50% withdrawal threshold was  
25 determined using the up-down method of Dixon (1980). Testing was started with the 2.04 g hair, in the middle of the series. Stimuli were always presented in a consecutive way, whether ascending or descending. In the absence of a paw withdrawal response to the initially selected hair, a stronger stimulus was presented; in the event of paw withdrawal, the next weaker stimulus was chosen. Optimal threshold calculation by this method

requires 6 responses in the immediate vicinity of the 50% threshold, and counting of these 6 responses began when the first change in response occurred, e.g. the threshold was first crossed. In cases where thresholds fell outside the range of stimuli, values of 15.14 (normal sensitivity) or 0.41 (maximally allodynic) were respectively assigned. The resulting pattern of positive and negative responses was tabulated using the convention, X = no withdrawal; O = withdrawal, and the 50% withdrawal threshold was interpolated using the formula:

$$50\% \text{ g threshold} = 10^{(X_f + k\delta)} / 10,000$$

10

where  $X_f$  = value of the last von Frey hair used (log units);  $k$  = tabular value (from Chaplan et al. (1994)) for the pattern of positive / negative responses; and  $\delta$  = mean difference between stimuli (log units). Here  $\delta = 0.224$ .

15 Von Frey thresholds were converted to percent of maximum possible effect (% MPE), according to Chaplan et al. 1994. The following equation was used to compute % MPE:

$$\% \text{ MPE} = \frac{\text{Drug treated threshold (g)} - \text{allodynia threshold (g)}}{\text{Control threshold (g)} - \text{allodynia threshold (g)}} \times 100$$

20


#### ADMINISTRATION OF TEST SUBSTANCE

Rats were injected (subcutaneously, intraperitoneally, or orally) with a test substance prior to von Frey testing, the time between administration of test compound and the von Frey test varied depending upon the nature of the test compound.

25

**CLAIMS**

1. A compound according to formula I  
formula I



wherein

**m and n** is each and independently an integer of from 0-3, and one or more of the  
10 hydrogens in such an alkylene-chain may optionally be substituted by anyone of  
**C<sub>1</sub> - C<sub>6</sub> alkyl, C<sub>1</sub> - C<sub>6</sub> alkoxy, or hydroxy; or**  
one or more of the methylene groups may optionally be substituted by a heteroatom such as  
**O, N or S;**

15 **R<sup>1</sup>** is selected from hydrogen, a branched or straight **C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkenyl, C<sub>3</sub>-C<sub>8</sub>**  
cycloalkyl, **C<sub>4</sub>-C<sub>8</sub>(alkyl-cycloalkyl)** wherein alkyl is **C<sub>1</sub>-C<sub>2</sub> alkyl** and cycloalkyl is **C<sub>3</sub>-C<sub>6</sub>**  
cycloalkyl;

**R<sup>2</sup>** is selected from any of

20

**(i) hydrogen;**

**(ii) a straight or branched C<sub>1</sub>-C<sub>6</sub> alkyl, said alkyl optionally comprising one or more**  
saturated bonds (double or triple bonds) within the alkyl chain;

25



(iii)  $-(\text{CH}_2)_q\text{-aryl}$ ];

(iv)  $-(\text{CH}_2)_r\text{-heteroaryl}$ ] where the heteroaryl has from 5 to 10 atoms and the heteroatom being selected from any of S, N and O;

and wherein the aryl and heteroaryl may optionally and independently be substituted by  
5 1 or 2 substituents Y where each Y is as defined below; and wherein q and r is each and independently an integer of from 0 to 3;

(v)  $\text{C}_3\text{-C}_{10}$  cycloalkyl, optionally comprising one or more unsaturations and optionally substituted by one or more heteroaryl(s) where the heteroaryl has from 5 to 10 atoms and  
10 the heteroatom being selected from any of S, N and O;

and wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents Y where each Y is as defined below;

(vi)  $\text{C}_6\text{-C}_{10}$  aryl, optionally and independently substituted by one or more heteroaryl(s)  
15 having from 5 to 10 atoms and the heteroatom(s) being selected from any of S, N and O and wherein the heteroaryl may optionally and independently be substituted by 1 or 2 substituents Y wherein each Y is as defined below;

(vii) heteroaryl having from 5 to 10 atoms and the heteroatom being selected from  
20 any of S, N and O; wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents Y wherein each Y is as defined below;

*or*

$\text{R}^1$  and  $\text{R}^2$  may optionally form a heterocyclic ring;

$R^3$  is selected from anyone of

(i) hydrogen;

5

(ii) a straight or branched  $C_1$ - $C_6$  alkyl, said alkyl optionally comprising one or more saturated bonds (double or triple bonds) within the alkyl chain;

(iii)  $-[(CH_2)_q\text{-aryl}]$  wherein  $q$  is an integer of from 0 to 3, and wherein the aryl may  
10 optionally be substituted by one or more heteroaryl(s) having from 5 to 10 atoms and the heteroatom being selected from any of S, N and O; and wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents Y wherein each Y is as defined below;

(iv) heteoaryl- $(C_5 - C_{10}$  alkyl), where the heteoaryl has from 5 to 10 atoms and the  
15 heteroatom being selected from any of S, N and O, and wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents Y where each Y is as defined below;

(v)  $C_3$ - $C_{10}$  cycloalkyl, optionally comprising one or more unsaturations and optionally  
20 substituted by one or more heteroaryl(s) having from 5 to 10 atoms and the heteroatom being selected from any of S, N and O, and wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents Y where each Y is as defined below;

25

(vi)  $-[(C_3$ - $C_6$  cycloalkyl)- $(CH_2)_q]$  wherein  $q$  is an integer of from 1 to 3;

$R^4$  is selected from

(i) hydrogen;

(ii) a straight or branched C<sub>1</sub>-C<sub>6</sub> alkyl, said alkyl optionally comprising one or more saturated bonds (double or triple bonds) within the alkyl chain;

5

(iii) -[(CH<sub>2</sub>)<sub>q</sub>-aryl] wherein q is an integer of from 0 to 3, and wherein the aryl may optionally be substituted by one or more heteroaryl(s) having from 5 to 10 atoms and the heteroatom being selected from any of S, N and O; and wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents Y wherein each Y is

10

(iv) heteroaryl-(C<sub>5</sub> - C<sub>10</sub> alkyl), where the heteroaryl has from 5 to 10 atoms and the heteroatom being selected from any of S, N and O, and wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents Y where each Y is as

15

(v) C<sub>3</sub>-C<sub>10</sub> cycloalkyl, optionally comprising one or more unsaturations and optionally substituted by one or more heteroaryl(s) where the heteroaryl has from 5 to 10 atoms and the heteroatom being selected from any of S, N and O;

20

and wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents Y where each Y is as defined below;

(vi) C<sub>6</sub>-C<sub>10</sub> aryl, optionally and independently substituted by one or more heteroaryl(s) having from 5 to 10 atoms and the heteroatom(s) being selected from any of S, N and O

25

and wherein the heteroaryl may optionally and independently be substituted by 1 or 2 substituents Y wherein each Y is as defined below;

(vii) heteroaryl having from 5 to 10 atoms and the heteroatom being selected from any of S, N and O; wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents Y wherein each Y is as defined below;

5

$R^5$  is selected from anyone of

(i) hydrogen;


10 (ii) a straight or branched  $C_1$ - $C_6$  alkyl, said alkyl optionally comprising one or more saturated bonds (double or triple bonds) within the alkyl chain;

(iii)  $-(CH_2)_q$ -aryl] wherein q is an integer of from 0 to 3, and wherein the aryl may optionally be substituted by one or more heteroaryl(s) having from 5 to 10 atoms and the  
15 heteroatom being selected from any of S, N and O; and wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents Y wherein each Y is as defined below;

(iv) heteroaryl- $(C_5 - C_{10}$  alkyl), where the heteroaryl has from 5 to 10 atoms and the  
20 heteroatom being selected from any of S, N and O, and wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents Y where each Y is as defined below;

(v) **C<sub>3</sub>-C<sub>10</sub> cycloalkyl**, optionally comprising one or more unsaturations and optionally substituted by one or more heteroaryl(s) having from 5 to 10 atoms and the heteroatom being selected from any of S, N and O, and wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents Y where each Y is as defined below;

(vi)



10

wherein

**R<sup>7</sup>, R<sup>8</sup>, R<sup>9</sup>, R<sup>10</sup> and R<sup>11</sup>** is each and independently selected from

15

(a) **hydrogen;**

(b) **a straight or branched C<sub>1</sub>-C<sub>6</sub> alkyl**, said alkyl optionally comprising one or more saturated bonds (double or triple bonds) within the alkyl chain;

(c)  $-(\text{CH}_2)_q\text{-aryl}$ ] wherein  $q$  is an integer of from 0 to 3, and wherein the aryl may optionally be substituted by one or more heteroaryl(s) having from 5 to 10 atoms and the heteroatom being selected from any of S, N and O; and wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents  $Y$  wherein each  $Y$  is as defined below;

(d) **heteoaryl-(C<sub>5</sub> - C<sub>10</sub> alkyl)**, where the heteroaryl has from 5 to 10 atoms and the heteroatom being selected from any of S, N and O, and wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents  $Y$  where each  $Y$  is as defined below;

(e) **C<sub>3</sub>-C<sub>10</sub> cycloalkyl**, optionally comprising one or more unsaturations and optionally substituted by one or more heteroaryl(s) where the heteroaryl has from 5 to 10 atoms and the heteroatom being selected from any of S, N and O; and wherein the aryl and heteroaryl may optionally and independently be substituted by 1 or 2 substituents  $Y$  where each  $Y$  is as defined below;

(f) **C<sub>6</sub>-C<sub>10</sub> aryl**, optionally and independently substituted by one or more heteroaryl(s) having from 5 to 10 atoms and the heteroatom(s) being selected from any of S, N and O and wherein the heteroaryl may optionally and independently be substituted by 1 or 2 substituents  $Y$  wherein each  $Y$  is as defined below;

or

**R<sup>4</sup> and R<sup>5</sup>** may optionally form a heterocyclic ring;

Y is each and independently selected from any of hydrogen, CH<sub>3</sub>; NO<sub>2</sub>;

—(CH<sub>2</sub>)<sub>p<sup>1</sup></sub>CF<sub>3</sub>; halogen; —CONR<sup>a</sup>R<sup>b</sup>; —COOR<sup>a</sup>; —COR<sup>a</sup>; —(CH<sub>2</sub>)<sub>p<sup>2</sup></sub>NR<sup>a</sup>R<sup>b</sup>;

—(CH<sub>2</sub>)<sub>p<sup>3</sup></sub>CH<sub>3</sub>(CH<sub>2</sub>)<sub>p<sup>4</sup></sub>SOR<sup>a</sup>R<sup>b</sup>; —(CH<sub>2</sub>)<sub>p<sup>5</sup></sub>SO<sub>2</sub>R<sup>a</sup>; —(CH<sub>2</sub>)<sub>p<sup>6</sup></sub>SO<sub>2</sub>NR<sup>a</sup>; -O-Aryl;

C<sub>4</sub>-C<sub>8</sub>(alkyl-cycloalkyl) wherein alkyl is C<sub>1</sub>-C<sub>2</sub> alkyl and cycloalkyl is C<sub>3</sub>-C<sub>6</sub>


cycloalkyl; and 1 or 2 heteroaryl(s) having from 5 to 10 atoms and the heteroatom(s) being selected from any of S, N and O; and wherein



R<sup>a</sup> and R<sup>b</sup> is each and independently selected from hydrogen, a branched or straight C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkenyl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl; and wherein

p<sup>1</sup>, p<sup>2</sup>, p<sup>3</sup>, p<sup>4</sup>, p<sup>5</sup> and p<sup>6</sup> is each and independently 0, 1 or 2.


as well as pharmaceutically acceptable salts, isomers, hydrates, isoforms and prodrugs thereof.

2. A compound according to claim 1, which compound is anyone selected from










5







1  
 2  
 3  
 4  
 5  
 6  
 7  
 8  
 9  
 10  
 11  
 12  
 13  
 14  
 15  
 16  
 17  
 18  
 19  
 20  
 21  
 22  
 23  
 24  
 25  
 26  
 27  
 28  
 29  
 30  
 31  
 32  
 33  
 34  
 35  
 36  
 37  
 38  
 39  
 40  
 41  
 42  
 43  
 44  
 45  
 46  
 47  
 48  
 49  
 50  
 51  
 52  
 53  
 54  
 55  
 56  
 57  
 58  
 59  
 60  
 61  
 62  
 63  
 64  
 65  
 66  
 67  
 68  
 69  
 70  
 71  
 72  
 73  
 74  
 75  
 76  
 77  
 78  
 79  
 80  
 81  
 82  
 83  
 84  
 85  
 86  
 87  
 88  
 89  
 90  
 91  
 92  
 93  
 94  
 95  
 96  
 97  
 98  
 99  
 100




3. A compound according to any of the preceding claims, in form of its hydrochloride, sulfate, tartrate or citrate salts.
4. A compound according to any of claims 1-3 for use in therapy.
5. A compound according to claim 3, wherein the therapy is pain management.
6. A compound according to claim 3, wherein the therapy is directed towards gastrointestinal disorders.
7. A compound according to claim 3, wherein the therapy is directed towards spinal injuries.
8. A compound according to claim 3, wherein the therapy is directed to disorders of the sympathetic nervous system.
9. Use of a compound according to formula (I) of claim 1 for the manufacture of a medicament for use in the treatment of pain.

10. Use of a compound according to formula (I) of claim 1 for the manufacture of a medicament for use in the treatment of gastrointestinal disorders.
11. Use of a compound according to formula (I) of claim 1 for the manufacture of a medicament for use in the treatment of spinal injuries.
12. A compound according to any of claims 1-3, further characterised in that it is isotopically labelled.
13. Use of a compound according to claim 12 as a diagnostic agent.
14. An isotopically labelled compound of the formula (I) of claim 1.
15. A diagnostic agent comprising a compound of the formula (I) of claim 1.
16. A pharmaceutical composition comprising a compound of the formula (I) according to claim 1 as an active ingredient, together with a pharmacologically and pharmaceutically acceptable carrier.

17. A process for the preparation of a compound of the formula (I) according to claim 1, whereby


5 A)

(i) bis-amino xylylene of the formula




is converted into mono-(diBoc)-guanidinomethyl of the formula (II)

10




which thereafter is reacted with an aldehyde, providing a secondary amine of the general formula IV

15




wherein  $R^3$  is as defined in formula I of claim 1;

(ii) **compound IV** is subjected to an urea formation, providing a compound of the formula  
5 (VI)



which finally is deprotected, providing a compound of the general formula VII




10

wherein


$R^1$ ,  $R^2$  and  $R^3$  are as defined in formula I of claim 1; or

B)


(i) a compound of the formula (VIII)




wherein X is CN and Y is CHO, is subjected to a reductive amination using a primary amine  $R^3NH_2$  wherein  $R^3$  is as defined in formula I of claim 1, providing a compound of the formula (IX)



wherein  $R^3$  is as defined in formula I of claim 1, which thereafter is subjected to an urea reaction by using a chloroformate of the formula (V)




which finally is reduced, providing a compound of the formula (XI)




; or

(ii) compound (VIII) wherein X is  $\text{CH}_2\text{Br}$  and Y is  $\text{CN}$ , is reacted with an amine  $\text{R}^4\text{R}^5\text{NH}$  wherein  $\text{R}^4$  and  $\text{R}^5$  are as defined in formula I of claim 1, providing a compound of the formula (XII)



which is reduced and reacted with an aldehyde  $\text{R}^3\text{CHO}$  wherein  $\text{R}^3$  is as defined in formula I of claim 1, and finally subjected to an urea action with a chloroformate (V) as in step (i), providing a compound of the formula (XV)



wherein  $\text{R}^1, \text{R}^2, \text{R}^3, \text{R}^4$  and  $\text{R}^5$  are as defined in formula I of claim 1.

5 18. A method for the treatment of pain, whereby an effective amount of a compound of the formula (I) according to claim 1 is administered to a subject in need of pain management.


10 19. A method for the treatment of gastrointestinal disorders, whereby an effective amount of a compound of the formula (I) according to claim 1, is administered to a subject suffering from said gastrointestinal disorder.

20. A method for the treatment of spinal injuries, whereby an effective amount of a compound of the formula (I) according to claim 1, is administered to a subject suffering from said spinal injury.



Abstract

Compounds of general formula I



are disclosed and claimed in the present application, as well as their pharmaceutically acceptable salts, pharmaceutical compositions comprising the novel compounds and their use in therapy, in particular in the management of pain.



2  
3